

GCSE Chemistry Complete Revision Summary



a) Atomic Structure and Mixtures

- b) Periodic Table
- c) Structure and Bonding
- d) Quantitative Chemistry
- e) Chemical Changes
- f) Energy Changes

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272

Chemicals of the natural environment

Ionic Bonding Nanoparticles

Covalent Bonding Graphere and Fullerene


Metallic Bonding

State of Matter

Ionic compounds

Covalent Compounds

Diamond and Graphite





IONIC BONDING: Metals and Non Metals

BitPaper Share Knowledge

www.expertguidance.co.uk

It is between a metal and a non metal

Metal gains an electron and become positively charged.

Non- Metal looses an electron and becomes negatively charged.

There is a strong electrostatic force of attraction between opposite charged ions resulting in ionic bonding.

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272




Al = 2, 8, 3F = 2, 7



Dot and Cross Diagram

Write the symbols
Write electronic configuration
show outer electrons
show transfer
show charges





IONIC BONDING: Metals and Non Metals



www.expertguidance.co.uk

It is between a metal and a non metal

1601L

Metal gains an electron and become positively charged.

Non- Metal leas an electron and becomes negatively charged.


There is a strong electrostatic force of attraction between opposite charged ions resulting in ionic bonding.

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272 Example: Magnesium Chloride

Mg=2,8,2 Cl, 2,8,7



Write the symbols
Write electronic configuration
show outer electrons
show transfer
show charges





Ionic Compound Properties

- Brittle solids with definite crystal shapes
- Good insulators in solid form, but become good conductors in liquid or dissolved form
- High melting and boiling point compared to molecular compounds

Source: Flickr.com

In lonic compounds, there is a strong electrostatic force of attraction between the opposite charged ions. This results in the formation of giant ionic lattice.

In the solid form, the ions are not free to move as they are held together by strong electrostatic force of attraction. In molten or when they are dissolved in water the ions are free to move and conduct electricity.

In Ionic compounds, there is a strong electrostatic force of attraction between the opposite charged ions. This results

in the formation of giant ionic lattice. Large amount of energy is required to overcome the strong electrostatic force of attraction. Therefore, ionic compounds have high melting and boiling point.

Greater the charge of an ionic lattice, stronger is the electrostatic force of attraction. Greater the melting and bp.


For ex Aluminium chloride > Magnesium chloride> sodium chloride


→ Write the Symbols

→ Write the charges

(Upto group the charge is same as the group number. After group 4 it is group number -8)

→ Criss Cross





b) Magnesium Nitride

c) Calcium Oxide = (a D

d) Sodium Sulphide = Na, S

e) Sodium Chloride

f) Magnesium chloride Magq

g) Aluminium Chloride

Mg2+13-

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272





Writing Formulas For Binary Ionic Compounds | Identify the metal and | non-metal | | Calcium oxide | i) Write the symbols | | O | ii) Write the charges | | O | iii) Cross over the charges | | From top to bottom |

Source: Flickr.com





Group- ↓Perio		2	3		4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	1 H																		2 He
2	3 Li	4 Be												5 B	6 C	7 N	8 0	9 F	10 Ne
3	11 Na	12 Mg												13 Al	14 Si	15 P	16 S	17 Cl	18 Ar
4	19 K	20 Ca	21 Sc		22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr
5	37 Rb	38 Sr	39 Y		40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe
6	55 Cs	56 Ba	57 La	*	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 TI	82 Pb	83 Bi	84 Po	85 At	86 Rn
7	87 Fr	88 Ra	89 Ac	*	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og
				*	58	59	60	61	62	63	64	65	66	67	68	69	70	71	
					Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Но	Er	Tm	Yb	Lu	
*				90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr		

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272

Source: Wikimedia Commons



Positive	Negative
Ammonium NH T	Hydroxide - 04 Carbonate - (03 Sulphate - So4 Nitrate - No3
	Phosphate - 704

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272

FORMULAE OF COMPLEX IONS



- a) Magnesium HYdroxide = $M_3(6N)_2$
- b) Aluminium Sulphate = $A_1(Sv_1)_2$
- c) Calcium Phosphate = ((Po4)
- d) Magnesium nitrate = $M_3(N_3)_2$
- e) Calcium Carbonate = (awa
- f) Sodium Phosphate





COVALENT BONDING

BitPaper Share Knowledge

It is between two non metals

It involves the sharing of electrons between two non metals.

More than one electron pair can also be shared resulting in the formation of single double and triple bonds.


HCC H




420


















N2






BitPaper
Share Knowledge


www.expertguidance.co.uk

02



0=0

(02



⁻⁻ (=-



Properties of Ionic and **Covalent Compounds** Covalent Crystals • S.I.G • Lower Melting Point / · High melting Points + boiling point **boiling Points** Poor conductors · Hard, Brittle Usually soluble in H2O . Don't usually dissolved in Electrolytes Flammable

Source: Flickr.com

www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272

Properties of Covalent Compounds !!!!!



Simple Molecule

Giant Covalent

Diamond

Graphite

They have weak intermolecular forces in them so have a lower melting and a boiling points

Silicon Dioxide



The intermolecular forces increases with increase in size as the surface area between the molecules increases.

Therefore, polymers which have covalent bonding between them have high melting and boiling point due to increase in chain length.



GIANT COVALENT STRUCTURES !!!!





www.expertguidance.co.uk

Substances which have huge netword of atoms joined together by covalent bonds form giant covalent structures.









DIAMOND	GRAPHITE
It is hard.	It is soft and greasy.
It is an insulator	It is a conductor
It has a high density.	It has a lower density than diamond.
Each carbon atom is covalently bonded to four other carbon atoms giving it a strong rigid structure.	Carbon atoms are bonded in the form of layer in the form of hexagons. No covelent bonding between the layers so they can slide past. Each carbon atom is bonded with three other carbon leaving the fourth electron has delocalized
No delocalised electrons present	It has delocalised electrons
Used in cutting or jewellery	It is used in pencil leads.



PROPERTIES OF GRAPHITE !!!!!!



Q1 Why graphite is soft and slippery?

In graphite, Carbon atoms are bonded in the form of layers in the form of hexagons. No covelent bonding between the layers so they can slide past each other. The layers have only weak intermolecular forces between them. By applying a little pressure then layers can easily slide past each other making graphite soft and slippery.

Insert an apostrophe in the correct place in the sentence below.


Pupils coats should be hung on the pegs.

Q2 Why graphite conduct electricity?

In graphite, Carbon atoms are bonded in the form of layer in the form of hexagons. No covelent bonding between the layers so they can slide past. Each carbon atom is bonded with three other carbon leaving the fourth electron has delocalized, These delocalized electrons are mobiles electrons which can move and conduct electricity.



Fullerene: Hollow shaped molecule having hexagonal rings like a bucky ball.




Also known as bucky ball or buckminsterfullerene.

Carbon can be in the form of pentagon or hexagon rings Used as catalyst, drug delievery and treating cancer.



Graphene: Layer of interlocking hexagonal rings like single sheet of graphite.



It is a better conductor than graphite, light and have low density.

Used in making computer chips and flexible electronic displays. www.expertguidance.co.uk

mahima.laroyia@expertguidance.co.uk +447448352272



CARBON NANOTUBES !!!!!



Cylinderical fullerene with the length greater than the diameter.


High tensile strength

Used in making reinforced composite materials

High electronic conductivity

1

used in electronic industry



Source: Wikipedia






Layered structure

It is between two metals.

There are fixed positive ions present in the sea of delocalised electrons.

There is strong electrostatic force of attraction between fixed positive ions and delocalized electrons resulting in metallic bonding.



Fixed positive ions

Sea of delocalised electrons www.expertguidance.co.uk

mahima.laroyia@expertguidance.co.uk Source: Wikimedia Commons

+447448352272

Metals are malleable

Malleable means that the metals can be hammered into any shape.

Metals have layered structure and layers can slide past each other by hammering giving metals different shapes.

Metals are ductile

Ductile means that the metals can be drawn into thin wires.

Metals have layered structure and layers can slide past each other by hammering giving metals a wire shape.

Metallic Bonding

Atoms in a metal are arranged in a regular manner and vibrate about fixed positions.

The outermost electrons move freely, forming a 'sea of electrons' enveloping the positive metal ions.

Metals are good conductors of electricity

Metals have delocalised electrons.

They are mobile and conduct electricity.

These mobile electrons or delocalised electrons conduct heat and electricity.

Source: Flickr.com

Metals have high melting and boiling points

There is strong electrostatic force of attraction between fixed positive ions and delocalized electrons.

Large amount of energy is required to overcome strong electrostatic force of attraction.

www.expertguidance.co.u mahima.laroyia@expertgu +447448352272






ALLOYS



Alloys are the mixture of metals with another metal or a non metal which make the metal stronger.



Layers distorted by another metal or a non metals

Example: Steel is the alloy of iron which is more strong and resistant to corrossion.

In metals the particles are arrranged in layers. There is a regular arrangement of fixed positive ions which can slide past each by applying pressure.


In alloys there is a mixture of metals with another metal or a non metals. Another metal being different in shape and size distort the regular arrangement of the metal lattice.


As a result the layers of the metal can no longer slide past each other making it strong

NANOPARTICLES




Nanoparticles are the particles that deals with the paricles of size 1 to 100 nm.





Volume = side X side x side





Surface Area = 6 x side X side

Volume = side X side x side

As the size decreases the surface area to volume ratio increases.

Therefore Nano particles being very small in size have large surface area to volume ratio making thew very useful in Science and Medicine.

Share Knowledge



APPLICATIONS OF NANOPARTICLES



To kill cancer and tuomour cells

MEDICINES

For drug delievery

+447448352272

Used in Sunscreen to block sunlight

COSMETICS

They have large surface area to volume ratio.

CATALYST

www.expertguidance.co.uk Used in small quantities mahima.laroyia@expertguidance.go.uk

Self cleaning window panes

HOUSEHOLD

Nano particles breaks dirty in the presence of sunshine which is washed away by water while raining.





Due to small size can cause difficulty in breathing



They can accumulate in the envrionment and cause air pollution




Due to their large surface area a small spark can result in violent explosion making them risky to use.



They are toxic and cause breathing and respiratory problems.



Due to their small size they can also cause water pollution and risk the aquation life.




			BitPaper Share Knowledge
SOLIDS	LIQUIDS	GASES	Thate knowledge
articles are close	Particles are slightly	Particles are far apart.]

closer to each other. to each other. Have fixed shape Do not have fixed shape Do not have fixed shape Very weak forces Strong forces Weak forces between the particles between the particles. between the particles Do not have fixed volume Have fixed volume Have definite volume cannot be Can be compressed Highly compressible compressed Cannot flow Can flow Can flow

Source: Flickr.com





*EXPERT GUIDANCE

KEY TERMS !!!!



www.expertguidance.co.uk

Ions — charged atoms with unequal number of protons and electrons.

Ionic Bonding — bond formed between a metal and a non metal which involves complete transfer of electrons from metal to a non metal.

Dot and Cross — diagram that show transfer of electron in an ionic bond or sharing of electrons in a covalent bond.

Covalent Bonding bonding between two non metals which involves sharing of electrons.

Metallic Bonding

bonding in metals which involves strong electrostatic forces of attraction between fixed positive ions and delocalised electrons.

Intermolecular Forces — The forces between the molecules which determines the melting or a boiling point.

Giant Covalent Molecules — Covalently bonded molecules which forms large giant structure.

Polymers — Molecules which are made up of many repeating units

Fullerene Allotrope of carbon which forms a cage like structure like bucky ball.

Graphene Allotrope of carbon which is equivalent to single layer of graphite

Alloys Mixture of metals with another metal or a non metal.

Nanoparticles - Particles which are of the size of 1 nm to 100 nm.

Nanoscience—It is the branch of science that deals with nanoparticles.

State of Matter-Different forms that a matter can take. They are solids, liquids and gas.

Solids — States of matter with fixed shape and volume.

Liquids— States of matter without fixed shape but fixed volume.

Gases States of matter with fixed shape and volume.

www.expertguidance.co.ul mahima.laroyia@expertgu +447448352272



TEST YOURSELF !!!!!



www.expertguidance.co.uk

Q3 Differentiate Between Diamond and Graphite

Q1 Name the type of bonding in the following compounds :-

- a) Sodium Chloride
- b) Magnesium
- c) Nitrogen
- d) Carbon Dioxide
- e) Water
- f) Ammonia

Q2 Draw dot and cross diagram to represent bonding in the following

- a) sodium chloride
- b) Water
- c) Magnesium

Q4 Why Ionic compounds do not conduct electricity in solids?

Q5 Why Alloys are stronger than metals

Q6 Why alumunium has a stronger melting point than sodium

Q7 What are nanoparticles? Write the properties and applications of nanoparticles

Q1 Name the type of bonding in the following compounds :-

- a) Sodium Chloride —— Ionic
- b) Magnesium —— Metallic
- c) Nitrogen —— Covalent
- d) Carbon Dioxide Covalent
- e) Water ____ Covalent
- f) Ammonia —— Covalent



Q3 Differentiate Between Diamond and Graphite

bonded to four other carbon atoms giving it a strong rigid structure.

Carbon atoms are bonded in the form of layer in the form of hexagons. No covelent bonding between the layers so they can slide past. Each carbon atom is bonded with three other carbon leaving the fourth electron has delocalized

No delocalised electrons present

Used in cutting or jewellery

DIAMOND


mas delocalised electrons


It is used in pencil leads.


GRAPHITE

Q2 Draw dot and cross diagram to represent bonding in the following

- a) sodium chloride
- b) Water
- c) Magnesium







www.expertguidance.co.uk mahima.laroyia@expertguidance.co.uk +447448352272

Share Knowledge



BitPaper Share Knowledge

www.expertguidance.co.uk

Q4 Why Ionic compounds do not conduct electricity in solids?

In solids, the ions are held together by strong electrostatic force of attraction in the giant ionic lattice. In molten state the ions are free to move therefore conduct electricity.

Q5 Why Alloys are stronger than metals

Alloys are the mixture of metals which distors the regular arrangement of metal as a result of which layers are not able to slide past each other making alloys stronger than metals.

Q6 Why alumunium has a stronger melting point than sodium

Aluminium has a greater charge. Due to greater charge of aluminium there is a stronger electrostatic forces of attraction between fixed positive ions and delocalised electrons. As a result aluminium has a greater melting point than sodium.

Q7 What are nanoparticles? Write the properties and applications of nanoparticles

Nanoparticles are the particles betwen the size of 1 to 100 nm.

Due to smaller size they have large surface area to volume ratio making them highly useful in medicine, catalysts, cosmetics and electronic industry.



NEXT STEP !!!!!





Check the specification



Do Exam Style Questions on this topic

•