

a) Atomic Structure and Mixtures

b) Periodic Table

c) Structure and Bonding

d) Quantitative Chemistry

e) Chemical Changes

f) Energy Changes

Group 7 (halogens)

4.1.2.1 The periodic table

4.1.2.2 Development of the periodic table

4.1.2.3 Metals and non-metals

4.1.2.4 Group 0

4.1.2.5 Group 1

4.1.2.6 Group 7

GROUP 7 : Halogens (Salt Forming)

Element	Symbol	Electronic Configuration	State at room temperature
Fluorine	9F^{19}	2, 7	Yellow Gas
Chlorine	17Cl^{35}	2, 8, 7	Green Gas and pale green in solution
Bromine	35Br^{80}	2, 8, 18, 7	Volatile brown liquid yellow in solution
Iodine	53I^{127}	2, 8, 18, 18, 7	Volatile purple solid brown in solution
Astatine	85At^{210}	2, 8, 18, 32, 18, 7	Radioactive

REACTIVITY
DECREASES
DUE TO
DECREASE
IN
ELECTRON
AFFINITY


- They are non metals
- They gain an electron to form -1 ions.
- They have low melting and boiling points
- Their melting point increases down the group due to increases in intermolecular forces.
- They are found in pairs and exist as diatomic molecules (X_2)
- They are poisonous and smelly
- Their reactivity increase down the group
- Their density increases down the group.
- They are poor conductors of heat and electricity

HALOGEN REACTION

X [F, Cl, Br, I]

REACTION WITH HYDROGEN


→ They react with hydrogen to form hydrogen halides.




→ Reactivity decreases down the group so fluorine and chlorine reacts explosively and bromine and iodine reacts at higher temperature in the presence of catalyst.

REACTION WITH METALS

→ They react with metals to form ionic compounds. In ionic compounds, halogens gain one electron from the metals to form -1 ions and attain noble gas electronic configurations.




- ★ The more reactive halogen displaces the less reactive halogen from its salt
- ★ As the reactivity decreases down the group, the halogen at the top can take the position of the halogen at the bottom in its compounds and will displace the less reactive halogen.



F
Cl
Br
I

Decreasing reactivity ↓



$\text{F}_2 \longrightarrow$ can displace all halogens
 $\text{Cl}_2 \longrightarrow$ can displace all halogen except fluorine
 $\text{Br}_2 \longrightarrow$ can displace only iodine

WHY REACTIVITY OF GROUP 7 DECREASES DOWN THE GROUP ?

→ The Reactivity of Group 7 decreases down the group as the electron affinity or tendency to gain the electron decreases down the group.

React. by electron

FACTORS AFFECTING TENDENCY TO GAIN AN ELECTRON

To gain an electron, smaller nuclear charge, smaller size and less shielding is required.

→ Nuclear Charge $+$

Smaller the size of the atom, greater will be the force of the nucleus as the electron will be closer to the nucleus.

→ SHIELDING

Less electrons and shells, smaller will be the shielding which will in turn increase the nuclear charge.

→ SIZE OF ATOMS

Greater the size of the atom, the outer electron will become further away from the nucleus resulting in decreases in nuclear charge

Down the group the atom size increases due to increase in number of electron shells. As a result the nuclear charge decreases.

The size of the atom also increases down the group which makes the nuclear charge weaker

The electron shells also increases which decreases the effective nuclear charge on the incoming electron.

Due to all these factors, the nuclear charge decreases which decreases the tendency of gaining electrons down the group of halogen making them less reactive.

NEXT STEP !!!

- ★ Check the Specification
- ★ Do Exam questions on this chapter

www.expertguidance.co.uk
mahima.laroyia@expertguidance.co.uk
+447448352272